

1359 - Lip Topper (100.00g)

Ingredient	Function	Percent	Wgt (g)	Wgt (oz)	Vol (tsp.)
Phase A					
Distilled Water (aqua)	Diluent	74.70	74.70	2.63	14.94
Glycerin (glycerin)	Humectant	1.50	1.50	0.05	0.30
Magnesium Aluminum Silicate (magnesium aluminum silicate)	Thickener	1.50	1.50	0.05	0.30
PVP (polyvinylpyrrolidone)	Film Former	1.30	1.30	0.05	0.26
Phase B					
Silky Talc Talc, Dimethicone	Texturizer	2.00	2.00	0.07	0.40
GelMaker EMU (sodium acrylate / sodium acryloyldimethyl taurate copolymer, isohexadecane, polysorbate 80)	Thickener, Emulsifier	1.00	1.00	0.04	0.20
Squalane (squalane)	Emollient	1.50	1.50	0.05	0.30
Caprylyl Glycol EHG (caprylyl glycol, ethylhexylglycerin)	Preservative	1.00	1.00	0.04	0.20
Phase C					
Mica Diamond Sparkle (mica [CI 77019], titanium dioxide [CI 77891])	Colorant	13.00	13.00	0.46	2.60
Phenoxyethanol SA (phenoxyethanol, sorbic acid, caprylyl glycol)	Preservative	0.50	0.50	0.02	0.10
Mica Interference Copper (mica [CI 77019], titanium dioxide [CI 77891], silica)	Colorant	2.00	2.00	0.07	0.40

Method

Combine phase A ingredients, heating to 50 C to help hydrate the Magnesium Aluminum Silicate. Stir thoroughly until PVP and Magnesium Aluminum Silicate is dispersed well and everything is homogenous. Cool to 40 C. Add phase B ingredients one by one in order, stirring well after each addition. Add phase C as desired. Viscosity can be adjusted with GelMaker EMU. Instead of interference copper another colored mica can be used if needed.

Properties

A beautiful, glittery lip topper to spruce up a lipstick or liquid lipstick or wear beautifully on it's own. Add lip balm on top to help seal in the glitter and prevent transfer.